

V-11294-A



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [R]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search

PubMed



for

v-11294-a



Limits

Preview/Index

History

Clipboard

Details

Note: Performing your original search, *v-11294-a*, in PubMed will retrieve 3 citations.

Display

AbstractPlus



Show

20



Sort by



Send to



All: 1

Review: 0



☐ 1: Br J Clin Pharmacol. 2002 Nov;54(5):478-84.



Links

Pharmacokinetic and pharmacodynamic profile following oral administration of the phosphodiesterase (PDE)4 inhibitor V11294A in healthy volunteers.

Gale DD, Landells LJ, Spina D, Miller AJ, Smith K, Nichols T, Rotshteyn Y, Tonelli A, Lacouture P, Burch RM, Page CP, O'Connor BJ.

Purdue Frederick Inc., Norwalk, CT, USA.

AIMS: To assess the pharmacokinetic and pharmacodynamic profile of the novel PDE4 inhibitor V11294A (3-(3-cyclopentyloxy-4-methoxybenzyl)-6-ethylamino-8-isopropyl-3H purine hydrochloride) in healthy male volunteers. **METHODS:** This was a double-blind, single dose, randomized crossover study in eight healthy volunteers who received a single oral, fasting dose of V11294A (300 mg) or placebo. Blood samples were taken before and 0.5, 1, 2, 2.5, 3, 4, 6, 9, 12, 18 and 24 h after oral dosing for determination of plasma concentrations of V11294A. Blood samples were also taken before and 3 and 24 h after dosing for the assessment of the effect of V11294A on mononuclear cell proliferation and tumour necrosis factor (TNF) release in whole blood. **RESULTS:** Following a single oral dose of 300 mg V11294A, plasma concentrations of V11294A and its active metabolite V10332 reached C_{max} (ng ml⁻¹; mean +/- s.d.; 1398 +/- 298, 1000 +/- 400, respectively) after 2.63 +/- 0.79 and 5.9 +/- 2.3 h, respectively. For V11294A and V10332, t_{1/2} were 9.7 +/- 3.9 and 9.5 +/- 1.7 h, and AUC(0, infinity) were 18100 +/- 6100 and 18600 +/- 8500 ng ml⁻¹ h, respectively. At 3 h dosing, plasma concentrations of V11294A and V10332 (3-(3-cyclopentyloxy-4-methoxybenzyl)-8-isopropyl-3H-purin-6-ylamine) were 1300 +/- 330 and 860 +/- 300 ng ml⁻¹, 7 and 3 times their in vitro IC₅₀s for inhibition of TNF release and proliferation, respectively. Treatment with V11294A resulted in a significant reduction of lipopolysaccharide (LPS)-induced TNF release at 3 h (P < 0.001) and at 24 h (P < 0.05) post ingestion. The amount of TNF released (pmol ml⁻¹) in response to a submaximal concentration of LPS (4 ng ml⁻¹) was not significantly altered following placebo treatment (before 681 +/- 68 vs 3 h postdose 773 +/- 109, P = 0.27).

Related Links

The new phosphodiesterase 4 inhibitor roflumilast is efficacious in exercise-induced asthma and leads to suppression of LPS-stimulated TNF-alpha ex vivo [Br J Clin Pharmacol. 2002]

Pharmacodynamic and pharmacokinetic profile of S 17092, a new orally active prolyl endopeptidase inhibitor, in elderly healthy volunteers. A phase I study. [Br J Clin Pharmacol. 2000]

Safety, tolerability and pharmacokinetics of oral S-3304, a novel matrix metalloproteinase inhibitor, in single and multiple dose escalation studies in healthy volunteers [Br J Clin Pharmacol Ther. 2005]

Pharmacokinetics and safety of escalating single and repeat oral doses of GW420867X, a novel non-nucleoside reverse transcriptase inhibitor. [Eur J Clin Pharmacol. 2001]

The novel phosphodiesterase 4 inhibitor, CI-1044, inhibits LPS-induced TNF-alpha production in whole blood [Br J Clin Pharmacol. 2005]

See all Related Articles...

In contrast, there was a significant reduction in the amount of TNF released following treatment with V11294A (before 778 +/- 87 vs 3 h postdose 566 +/- 72, $P = 0.02$). Phytohaemagglutinin (PHA) stimulated the incorporation of [3H]-thymidine in whole blood prior to drug administration. V11294A inhibited the PHA-induced proliferation at 3 h ($P < 0.05$). No adverse reactions were noted following single oral administration of V11294A. **CONCLUSIONS:** A single oral 300 mg dose of V11294A administered to healthy volunteers results in plasma concentrations adequate to inhibit activation of inflammatory cells ex vivo, which persists for at least 24 h without any adverse reactions.

PMID: 12445026 [PubMed - indexed for MEDLINE]

Display ☒ AbstractPlus ☒ Show 20 ☒ Sort by ☒ Send to ☒

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 6 2007 07:35:03

AWD 12-281



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[Sign In] [R]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Search PubMed



for awd 12-281

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Note: Performing your original search, *awd 12-281*, in PubMed will retrieve 11 citations.

Display AbstractPlus



Show

20



Sort by



Send to



All: 1

Review: 0



☐ 1: J Pharm Pharmacol. 2003 Aug;55(8):1107-14.



Links

AWD 12-281, a highly selective phosphodiesterase 4 inhibitor, is effective in the prevention and treatment of inflammatory reactions in a model of allergic dermatitis.

Baumer W, Gorr G, Hoppmann J, Ehinger AM, Rundfeldt C, Kietzmann M.

Department of Pharmacology, Toxicology and Pharmacy, School of Veterinary Medicine, Buenteweg 17, D-30559 Hannover, Germany.
wolfgang.baeumer@tiho-hannover.de

AWD 12-281 (N-(3,5-dichloro-4-pyridinyl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxoacetamide), a phosphodiesterase 4 inhibitor, which is optimized for topical administration, was tested in a model of allergic dermatitis in mice. To obtain an allergic dermatitis, BALB/c mice were sensitized to toluene-2,4-diisocyanate (TDI). The allergic reaction was challenged by topical administration of TDI onto the mice ears. AWD 12-281 was tested for its anti-inflammatory potential by oral, intraperitoneal and topical administration. The phosphodiesterase 4 inhibitor, cilomilast (SB 207499), and/or the corticosteroid, diflorasone diacetate, were used as reference compounds. Given orally and intraperitoneally 2 h before as well as 5 and 24 h after TDI challenge, AWD 12-281 showed no, or only a transient inhibition of the allergen-induced ear swelling, whereas cilomilast significantly inhibited this ear swelling. Applied topically onto the ears before TDI challenge, AWD 12-281, cilomilast and diflorasone diacetate caused total inhibition of ear swelling 24 h after challenge, confirmed by a decrease of the pro-inflammatory cytokines interleukin-4, interleukin-6 and macrophage inhibitory protein-2. Administered topically after TDI challenge as therapeutic intervention, AWD 12-281 and diflorasone diacetate caused significant inhibition of ear swelling; cilomilast failed to do so. These results indicate that topically administered AWD 12-281 may be potent in the prevention and treatment of allergic/inflammatory skin diseases.

PMID: 12956900 [PubMed - indexed for MEDLINE]

Related Links

Effects of the phosphodiesterase 4 inhibitors SB 207499 and AWD 12-281 on the inflammatory reaction in a model of allergic dermatitis [J Pharm Pharmacol. 2003]

The phosphodiesterase 4 inhibitor AWD 12-281 is active in a new guinea-pig model of allergic skin inflammation predictive of human skin penetration and suppresses both Th1 and Th2 cytokines in mice. [J Pharm Pharmacol. 2005]

In vivo efficacy in airway disease models of N-(3,5-dichloropyrid-4-yl)-[1-(4-fluorobenzyl)-5-hydroxy-indole-3-yl]-glyoxylic acid amide (AWD 12-281), a selective phosphodiesterase 4 inhibitor for inhaled asthma [Pharmacol Exp Ther. 2003]

TARC and RANTES, but not CTACK, are induced in two models of allergic contact dermatitis. Effects of cilomilast and diflorasone diacetate on T-cell-attracting chemokines. [Br J Dermatol. 2004]

Anti-inflammatory potential of the selective phosphodiesterase 4 inhibitor N-(3,5-dichloro-pyrid-4-yl)-[1-(4-fluorobenzyl)-5-hydroxy-indole-3-yl]-glyoxylic acid amide (AWD 12-281), in human cell preparation [J Pharm Pharmacol Exp Ther. 2004]

See all Related Articles...

Display AbstractPlus



Show

20



Sort by



Send to



[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Mar 6 2007 07:35:03